Nitrogen Compounds

Question Paper 7

Level	International A Level
Subject	Chemistry
Exam Board	CIE
Topic	Nitrogen Compounds
Sub-Topic	
Paper Type	Theory
Booklet	Question Paper 7

Time Allowed: 80 minutes

Score: /66

Percentage: /100

Grade Boundaries:

A*	Α	В	С	D	Е	U
>85%	777.5%	70%	62.5%	57.5%	45%	<45%

For more awesome GCSE and A level resources, visit us at www.savemyexams.co.uk/

1 (a) Organohalogen compounds can undergo hydrolysis.

$$R-Cl + H_2O \rightarrow R-OH + HCl$$

State the relative rates of hydrolysis of the following compounds.

Explain your answer.

.....

(b) Aminolaevulinic acid is involved in the synthesis of haemoglobin and chlorophyll.

$$H_2N$$
 OH

aminolaevulinic acid

Name the **three** functional groups in aminolaevulinic acid.


.....[2

- **(c)** Aminolaevulinic acid reacts readily with bromoethane.
 - (i) Show the mechanism of the **first step** of this reaction on the diagram. Include all necessary curly arrows, lone pairs and relevant dipoles.

For more awesome GCSE and A level resources, visit us at www.savemyexams.co.uk/

(ii)	Name the mechanism in (c)(i).
(iii)	Identify the non-organic product formed in this reaction.
	[5]

(d) Three reactions of aminolaevulinic acid are shown. Draw the structures of the products W, X and Y in the boxes below.



(e) Aminolaevulinic acid can undergo polymerisation.

Draw the structure of the polymer showing **two** repeat units. The linkages between the monomer units should be shown fully displayed.

2	(a)	A r	nixture of amino acids can be separated by electrophoresis. During an electrophore	resis
		exp	periment,	
		•	different amino acids move in different directions, different amino acids move at different speeds, some amino acids do not move at all.	
		Exp	plain these observations.	
	(b)	(i)	A mixture of amino acids can also be separated by thin-layer chromatography. Identify the mobile and the stationary phases in this type of chromatography.	
			mobile phase	
			stationary phase	
		(ii)	What is the process by which thin-layer chromatography can separate a mixture?	
				[3]
	(c)	Sta	te three structural features of DNA.	
				. [3]

For more awesome GCSE and A level resources, visit us at www.savemyexams.co.uk/

(d) Some diseases are caused by a mutation in the DNA base sequence which results in one amino acid being replaced by another during protein synthesis. Suggest what changes in the interactions that form the tertiary structure would result from a mutation that replaced a valine residue with a serine residue.

	replaced by	$ \begin{array}{c} $	
val		ser	

[Total: 11]

For more awesome GCSE and A level resources, visit us at <u>www.savemyexams.co.uk/</u>

3	There are two important polymerisations that occur within living organisms – protein synthesis
	and the formation of DNA.

(a) Complete the table by placing a tick (\checkmark) in the correct column to indicate in which process each substance could be used.

substance	protein synthesis	formation of DNA
cysteine		
cytosine		
glutamine		
guanine		

[3]

(b)	DN.	A consists of a double helical structure.
	(i)	Describe the bonding between the two strands in DNA and state which part of each strand is joined by it.
	(ii)	How does the strength of this bonding relate to the mechanism of the replication of DNA?
		[4]
(c)		ne diseases are caused by changes in the structure of proteins. Explain the genetic is of these changes.
		[3]

[Total: 10]

another	in the	Periodic Table.		
(a) In t	he box	tes below, draw the 'dot-and- ds. Show outer electrons only. shape of each molecule.	cross' diagram of a molec	ule of each of these
		NH ₃	CH ₄	
		shape	shape	
	npound	is polar whereas methane is ds are different. in, using ammonia as the exam		
(ii)	Expla	in why the ammonia molecule	is polar.	
(iii)	State	one physical property of amm	onia which is caused by its	s polarity.

.....

(c) When ammonia gas is mixed with hydrogen chloride, white, solid ammonium ch formed.	nloride is
State each type of bond that is present in one formula unit of ammonium chlo how many of each type are present. You may draw diagrams.	ride and
	otal: 10]

For more awesome GCSE and A level resources, visit us at www.savemyexams.co.uk/

5 (a) Enzymes are particular types of proteins that catalyse chemical reactions. The efficiency of enzymes can be reduced by the presence of other substances known as inhibitors.


` '	State one example of a substance that can act as a <i>non-competitive</i> inhibitor in enzyme reactions.
	•

.....

.....

(ii)	For the inhibitor you have identified, explain why it is a non-competitive inhibitor.

(iii) The graph shows the rate of an enzyme-catalysed reaction against the substrate concentration in the absence of an inhibitor.



On the same axes, sketch a graph showing the rate of this reaction if a *competitive inhibitor* was present.

For more awesome GCSE and A level resources, visit us at www.savemyexams.co.uk/

(b) DNA is responsible for encoding the amino acid sequence to produce proteins.

Ribosome, tRNA and mRNA are all involved in the process of protein synthesis.

(i) Write ribosome, tRNA and mRNA in the boxes below to show the correct sequence in which they are involved.

DNA —	protein
(ii)	Sequences of three bases code for specific amino acids. The code UGA however does not usually code for an amino acid. Suggest its use.
	[3]
` '	ich of the energy used in biochemical reactions is provided by the hydrolysis of the blecule ATP.
(i)	What are the breakdown products of the hydrolysis of ATP?
(ii)	Give two uses for the energy released by ATP hydrolysis in cells.
	1
	2

[Total: 10]

[3]

6	In key r	eactions responsible for growth and repair in the human body, amino acids react
	togethe	r to form polymers known as proteins.
	(a)	What type of reaction is this polymerisation?
	(ii)	From stocks of glycine and alanine, it is possible to make the dipeptide gly-ala. Using the same three-letter abbreviations for the amino acids, give the structures of all other possible dipeptides that can be made from these stocks of amino acids.
		[3]
	(b)	DNA consists of a double helix formed by two strands held together by hydrogen bonds between base pairs. Sketch a section of DNA showing two base pairs, using blocks for the various components. You should label all of the components.
	(ii)	Suggest what the effect on DNA replication would be if the hydrogen bonds between the strands were replaced by stronger bonds, e.g. covalent bonds.
		[4]

(c)		ne diseases, such as sickle-cell anaemia, are caused by mutation resulting in a nge in the triplet code.
	(i)	Explain why some changes in the triplet code do not result in a change in the primary structure of a protein.
	(ii)	Suggest what change in the tertiary structure of a protein would result from a mutation that replaced aspartic acid with glycine.
		Suggest what change in the tertiary structure of a protein would result from a mutation that replaced aspartic acid with glycine. O H ₂ N—CH—C—OH CH ₂ CH ₂ C=O OH
		aspartic acid glycine
	(iii)	Sometimes a mutation can result in the <i>deletion</i> of a single base in DNA (or RNA) Explain why this is likely to have more serious consequences for the protein than the <i>replacement</i> of one base by another.
		[3
		[Total: 10