

# States of Matter

## Question Paper 4

<b>Level</b>	International A Level
<b>Subject</b>	Chemistry
<b>Exam Board</b>	CIE
<b>Topic</b>	States of Matter
<b>Sub-Topic</b>	
<b>Paper Type</b>	Theory
<b>Booklet</b>	Question Paper 4

**Time Allowed:** 51 minutes

**Score:** /42

**Percentage:** /100

**Grade Boundaries:**

A*	A	B	C	D	E	U
>85%	77.5%	70%	62.5%	57.5%	45%	<45%

1 Propane,  $C_3H_8$ , and butane,  $C_4H_{10}$ , are components of Liquefied Petroleum Gas (LPG) which is widely used as a fuel for domestic cooking and heating.

(a) (i) To which class of compounds do these two hydrocarbons belong?

.....

(ii) Write a balanced equation for the complete combustion of butane.

..... [2]

(b) When propane or butane is used in cooking, the saucepan may become covered by a solid black deposit.

(i) What is the chemical name for this black solid?

.....

(ii) Write a balanced equation for its formation from butane.

..... [2]

(c) Propane and butane have different values of standard enthalpy change of combustion.

Define the term *standard enthalpy change of combustion*.

.....  
.....  
..... [2]

(d) A  $125\text{ cm}^3$  sample of propane gas, measured at  $20\text{ }^\circ\text{C}$  and  $101\text{ kPa}$ , was completely burnt in air.

The heat produced raised the temperature of  $200\text{ g}$  of water by  $13.8\text{ }^\circ\text{C}$ .

Assume no heat losses occurred during this experiment.

(i) Use the equation  $pV = nRT$  to calculate the mass of propane used.

- (ii) Use relevant data from the *Data Booklet* to calculate the amount of heat released in this experiment.
- (iii) Use the data above and your answers to (i) and (ii) to calculate the energy produced by the burning of 1 mol of propane.

[5]

- (e) The boiling points of methane, ethane, propane, and butane are given below.

compound	CH <sub>4</sub>	CH <sub>3</sub> CH <sub>3</sub>	CH <sub>3</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>3</sub> (CH <sub>2</sub> ) <sub>2</sub> CH <sub>3</sub>
boiling point/K	112	185	231	273

- (i) Suggest an explanation for the increase in boiling points from methane to butane.
- .....
- .....
- .....
- (ii) The isomer of butane, 2-methylpropane, (CH<sub>3</sub>)<sub>3</sub>CH, has a boiling point of 261 K. Suggest an explanation for the difference between this value and that for butane in the table above.

.....

.....

.....

[4]

[Total: 15]

- 2 Methanol, CH<sub>3</sub>OH, is considered to be a possible alternative to fossil fuels, particularly for use in vehicles.

Methanol can be produced from fossil fuels and from agricultural waste. It can also be synthesised from carbon dioxide and hydrogen.

- (a) Define, with the aid of an equation which includes state symbols, the standard enthalpy change of formation of carbon dioxide.

equation .....

definition .....

.....


..... [3]

- (b) Relevant  $\Delta H_f^\ominus$  values for the reaction that synthesises methanol are given in th

compound	$\Delta H_f^\ominus / \text{kJ mol}^{-1}$
CO <sub>2</sub> (g)	-394
CH <sub>3</sub> OH(g)	-201
H <sub>2</sub> O(g)	-242

- (i) Use these values to calculate  $\Delta H_{\text{reaction}}^\ominus$  fo

Include a sign in your answer.



$$\Delta H_{\text{reaction}}^\ominus = \dots\dots\dots \text{kJ mol}^{-1}$$

- (ii) Suggest **one** possible environmental advantage of this reaction. Explain your answer.

.....

.....

[5]

- (c) The synthesis of methanol is carried out at about 500K with a pressure of between 40 and 100 atmospheres (between  $4 \times 10^6$  Pa and  $10 \times 10^7$  Pa) and using a catalyst. The use of such conditions will affect both the rate of reaction and the equilibrium yield.

In the spaces below, explain the effects of higher temperature, higher pressure, and the use of a catalyst on the **equilibrium yield** of methanol.

**higher temperature**

effect .....

explanation .....

.....

**higher pressure**

effect .....

explanation .....

.....

**use of catalyst**

effect .....

explanation .....

.....

[6]

[Total: 14]

- 3 The elements carbon and silicon are both in Group IV of the Periodic Table. Carbon is the second most abundant element by mass in the human body and silicon is the second most common element in the Earth's crust.

Carbon and silicon each form an oxide of general formula  $XO_2$ .

At room temperature,  $CO_2$  is a gas while  $SiO_2$  is a solid with a high melting point.

- (a) Briefly explain, in terms of the chemical bonds and intermolecular forces present in **each** compound, why  $CO_2$  is a gas and  $SiO_2$  is a solid at room temperature.

.....  
.....  
.....  
..... [3]

- (b) Draw a simple diagram to show the structure of  $SiO_2$ . Your diagram should contain at least **two** silicon atoms **and** show clearly how many bonds each atom forms.

[2]

CO<sub>2</sub> does not behave as an ideal gas.

(c) (i) State the basic assumptions of the kinetic theory as applied to an ideal gas.

.....  
.....  
.....  
.....  
.....  
.....

(ii) Suggest **one** reason why CO<sub>2</sub> does not behave as an ideal gas.

..... [5]

Carbon exists in a number of forms, one of which is a conductor of electricity and one of which is a non-conductor of electricity. Silicon is the main component of most semi-conductors.

(d) Graphite is the form of carbon that is a conductor of electricity. Give a simple explanation for this property.

..... [1]

When carbon and silicon(IV) oxide are heated together at about 2000 °C, silicon carbide, SiC, is formed. Silicon carbide is a hard material which is widely used as an abrasive and in ceramics.

(e) (i) Construct an equation for the reaction of carbon and silicon(IV) oxide.

.....

(ii) SiC has a similar structure to one of the common forms of carbon. Which form is this? Give a reason for your answer.

form .....

reason ..... [2]

[Total: 13]